


BETTER PREVENTION PRACTITIONER TRAINING INSTITUTE

IMMUNOSUPPRESSION TIP SHEET

Immunocompromised conditions include:

- HIV
- Undergoing cancer treatment (chemotherapy, radiation)
- Post-organ transplant therapy
- Medication use (chronic current use):



If the patient flags for any of the above conditions or treatments, refer to the primary care provider to determine whether increased screening is required.

Please note: this is not a complete list of medications or conditions which can cause immunosuppression. The medications listed refer to injectable or oral administration, not creams or topical treatment. Some medications are dose dependent. If you are not sure, consult your team.



IMMUNOSUPPRESSION TIP SHEET

ANTIMETABOLITES

Azathioprine (Imuran)
Methotrexate (Metoject,
Rheumatrex, Methotrexate Sodium)
Purinethol (Mercaptopurine)

Betamethasone
Budesonide (Cortiment, Entocort)
Hydrocortisone
Methylprednisone
Prednisone
Prednisolone

CORTICOSTEROIDS

TNF Blockers

Adalimumab (Humira)
Baricitinib (Olumiant)
Certolizumab pegol (Cimzia)
Etanercept (Enbrel)
Infliximab (Remicade)

Apremilast (Otezla)
Basiliximab (Simulect)
Belimumab (Benlysta)
Cyclosporine (Neoral)
Mycophenolate (Cell Cept, Myfortic)
Tacrolimus (PROgraf, ADVagraf)
Sirolimus (Rapamune)
Sulfasalazine (Azulfidine)

OTHER